


Journal of Diagnostics and Treatment of Oral and Maxillofacial Pathology

7 2025



The cover images indicate the key direction of the journal and highlight an article titled “The ‘Beveled One-and-a-Half-Barrel’ Fibula Transplant with Virtual Surgical Planning and CT-Guided Implant Surgery for Prosthetic Rehabilitation in Posterior Mandible Defects: A Pictorial Essay” by Olindo MASSARELLI and Silvio Mario MELONI. The article was published in Volume 6, Issue 3.

This is a monthly open access and peer-reviewed publication. This is the first Ukrainian journal dedicated exclusively to oral and maxillofacial surgery, registered in 2016. The Editor in Chief is Oleksii Tymofeiev, Kyiv, Ukraine.



TANTUM VERDE®

QUICK RELIEF FROM PAIN
AND INFLAMMATION IN THE
MOUTH AND THROAT¹

**AN INTEGRAL COMPONENT OF THE TREATMENT
OF PAIN AND INFLAMMATION IN THE ORAL CAVITY
IN 60 COUNTRIES WORLDWIDE!²**



Reg. № UA/3920/01/01

**LOCAL ANESTHETIC
AND ANTI-INFLAMMATORY
EFFECT¹**

- **JAWS
FRACTURES³**
- **IMPLANTS
PLACEMENT⁴**
- **WOUNDS OF ORAL
CAVITY⁵**



SUMMARY OF PRODUCT CHARACTERISTICS

NAME OF THE MEDICINAL PRODUCT. Tantum Verde 0.15% mouthwash. **QUALITATIVE AND QUANTITATIVE COMPOSITION.** Each 100 ml contains: active ingredient: benzydamine hydrochloride 0.15 g (equivalent to 0.134 g of benzydamine). **Therapeutic indications.** Treatment of symptoms such as irritation/inflammation including those associated with pain in the oropharyngeal cavity (e.g. gingivitis, stomatitis and pharyngitis), including those resulting from conservative or extractive dental therapy. **Posology and method of administration.** Pour 15 ml of Tantum Verde mouthwash into the measuring cup, 2-3 times per day, using it either at full concentration or diluted. If diluted, add 15 ml of water to the graduated cup. Do not exceed the recommended dosage. **Contraindications.** Hypersensitivity to benzydamine or to any of the excipient. **PHARMACOLOGICAL PROPERTIES. Pharmacodynamic properties.** Pharmacotherapeutic group: Stomatologic drugs: other agents for local oral treatment, ATC code: A01AD02. Clinical studies demonstrate that benzydamine is effective in relieving suffering from localised irritation of the mouth and pharynx. In addition, benzydamine possesses a moderate local anaesthetic effect. **Pharmacokinetic properties. Absorption.** Absorption through the oropharyngeal mucosa is demonstrated by the presence of measurable quantities of benzydamine in human plasma. These levels are insufficient to produce systemic effects. **Distribution.** When applied locally, benzydamine has been shown to accumulate in inflamed tissues where it reaches effective concentrations because of its capacity to penetrate the epithelial lining.

Information about medicines. Information for health care professionals for use in professional activities.

1. Інструкція для медичного застосування лікарського засобу Тантум Верде®, розчин для ротової порожнини, РП № UA/3920/01/01, затверджено Наказом Міністерства охорони здоров'я України № 636 від 01.10.2015.

2. <http://www.angelini-pharma.com/wps/wcm/connect/com/home/Angelini+Pharma+in+the+world/>

3. Тимофеев А.А. и др. "Особенности гигиены полости рта для профилактики воспалительных осложнений при переломах нижней челюсти". Современная стоматология 2015;1(75):52-8.

4, 4.5. Tymofiev O.O. et al "Prevention of inflammatory complications upon surgeries in maxillofacial region". J Diagn Treat Oral Maxillofac Pathol. 2017;1:105-12.

Clinical and CT images are courtesy of: Ievgen Fesenko (Department of Oral & Maxillofacial Surgery, PHEI "Kyiv Medical University", Kyiv, Ukraine), Oleg Mostakov ("SCIEDECE—Scientific Center of Dentistry & Ultrasound Surgery" Kyiv, Ukraine).



04119, Kiev, Melnikova str. 83D, of. 404.
Tel.: (044) 538-01-26
Fax: (044) 538-01-27



About the Journal

JULY 2025 • VOLUME 9 • ISSUE 7
www.djournal.org

Official Title

Journal of Diagnostics and Treatment of Oral and Maxillofacial Pathology

Official Title in Ukrainian

Журнал діагностики та лікування оральної і щелепно-лицевої патології

Standard Abbreviation: ISO 4

J. Diagn. Treat. Oral Maxillofac. Pathol.

Acronym

JDTOMP

International Standard Serial Number (ISSN)

ISSN 2522-1965 (online)

Aims & Scope

This is a monthly open access and peer-reviewed oral and maxillofacial surgeons. The journal is focused on trauma, microvascular and jaw reconstructive surgery, dental implants, salivary gland tumors/diseases, TMJ lesions, virtual surgical planning, implementation of ultrasonography into the practice of oral and maxillofacial surgeons.

Editorial Board (EB) Composition

- EB shows significant geographic diversity representing 36 opinion leaders from 15 countries: Brazil, Canada, Colombia, Greece, Hong Kong (SAR), India, Indonesia, Israel, Italy, Slovak Republic, Spain, Ukraine, United Arab Emirates, United Kingdom, and United States.
- The majority of the EB Members have a discernible publication history in Scopus, Web of Science, and journals with a high impact factor.
- The publication records of all EB members are consistent with the stated scope and published content of the journal.
- The journal has several full-time professional editors.
- Gender distribution of the editors: 11.11% women (4 persons), 88.89% men (32 persons), 0% non-binary/other, and 0% prefer not to disclose.

Frequency

12 issues a year (from January 2020)

Publishing Model

The *Journal of Diagnostics and Treatment of Oral and Maxillofacial Pathology* is a fully online-only open access and peer-reviewed publication.

Open Access

This journal is licensed under the CC BY-NC-SA license.

Types of Peer Review

The journal employs “double blind” and open reviewing.

Article Publishing Charge (APC)

The APC in this journal is 519 Euro (\$590 USD) and 259 Euro (\$295 USD)(excluding taxes) depending on the article's type. Details at website: www.djournal.org.

Types of Articles Currently Published by the Journal

Editorials/Guest Editorials/Post Scriptum Editorials, Images, Case Reports/Case Series, Original Articles, Review Articles, Discussions, Paper Scans (*synonyms*: Review of Articles, Literature Scan), Book Scans (*synonym*: Book Reviews), Letters to the Editor (*synonym*: Letters), and Viewpoints.

State Registration in the Ministry of Justice of Ukraine

Registration: Jul 28, 2016 (Certificate: KB № 22251-12151 P)

Re-registration: May 21, 2019 (Certificate: KB № 23999-13839 ПП)

Re-registration: Aug 10, 2021 (Certificate: KB № 24951-14891 ПП).

State Registration in the National Council of Television and Radio Broadcasting of Ukraine

- The journal is registered as print media (Decision of 2024, April 11, No. 1225). The identifier of print media R30-04318.
- The journal is registered as online media (Decision of 2024, April 25, No. 1454). The identifier of online media R40-04708.

Journal is Included in

Encyclopedia of Modern Ukraine, Google Scholar, National Repository of Academic Texts, Register of Scientific Publications of Ukraine (also known as Ukrainian Scientific Periodicals or Register of Scientific Professional Publications of Ukraine), ResearchGate, Scilit, and Vernadsky National Library of Ukraine

Co-Founders

1. Shupyk National Healthcare University of Ukraine (formerly known as Shupyk National Medical Academy of Postgraduate Education).
2. Private Higher Educational Establishment “Kyiv Medical University.”
3. OMF Publishing, Limited Liability Company.

Publisher

OMF Publishing, LLC is an academic publishing company.

Address: 13-A Simferopolska Street, office 121, Kyiv 02096, Ukraine.

ROR ID: <https://ror.org/017s0zm38>.

E-mail: office@omfpublishing.com.

Website: www.omfpublishing.com.

Crossref Membership

OMF Publishing, LLC is a member of Publishers International Linking Association, Inc. which doing business as a Crossref. OMF Publishing's active membership: From February 2017 to present.

Digital Object Identifier (DOI) of the Journal

10.23999/j.djomp

Official Journal of the Association

Ukrainian Association for Maxillofacial and Oral Surgeons

Ukrainian Association for Maxillofacial and Oral Surgeons (UAMOS)

Address: 4-A Profesora Pidvysotskoho Street, Kyiv 01103, Ukraine.
Tel., fax: +38 044 528 35 17.

Website: www.uamos.org.

© 2025 OMF PUBLISHING, LLC

Editorial Board

JULY 2025 • VOLUME 9 • ISSUE 7
www.djournal.org

Editor in Chief

Oleksii O. Tymofieiev, ScD
Kyiv, Ukraine

Honorary Deputy Editor in Chief

Rui P. Fernandes, MD, DMD, FACS, FRCS(Ed)
Jacksonville, Florida, United States

Deputy Editors in Chief

Evangelos G. Kilipiris, DMD
Bratislava, Slovak Republic

John M. Le, MD, DDS
Jacksonville, Florida, United States

Camilo Mosquera, DDS
Galveston, Texas, United States

Section Editors

Bone Augmentation Techniques
Nardy Casap, MD, DMD
Jerusalem, Israel

Business
Ivan V. Nagorniak, MSc, PhD
Kyiv, Ukraine

Craniofacial Deformities
Sunil Richardson, MDS
Nagercoil, Tamil Nadu, India

Facial Feminization Surgery
Kyle Keojampa, MD, FACS
Los Angeles, California, United States

Facial Plastic Surgery
Tirbod Fattahi, MD, DDS, FACS
Jacksonville, Florida, United States

Head & Neck Oncologic Surgery
Todd C. Hanna, MD, DDS, FACS
New York, New York, United States

Head & Neck Radiology
Anil T. Ahuja, MBBS, MD, FRCR, FHKCR, FHKAM
Hong Kong, SAR

Images
Camilo Mosquera, DDS
Galveston, Texas, United States

Orthognathic Surgery
Mario Brinhole
São Paulo, São Paulo, Brazil

Robotic Surgery
Salam O. Salman, MD, DDS, FACS
Jacksonville, Florida, United States

TMJ Lesions/Disorders
Belmiro C. Vasconcelos, DDS, PhD
Recife, Pernambuco, Brazil

Trauma
Paul Coulthard, BDS, MDS, PhD, MFGDP(UK), FDSRCS(Eng), FDSRCS(OS), FDSRCPS(Glas), FFDTRCS(Ed), FDSRCS(Ed), FCGDent
London, United Kingdom

Editorial Board Members

Oleh M. Antonyshyn, MD, FRCS(C)
Toronto, Ontario, Canada

Anthony M. Bunnell, MD, DMD, FACS
Jacksonville, Florida, United States

Andi S. Budihardja, PhD
Lippo Village, Tangerang, Banten, Indonesia

Yurii V. Chepurnyi, ScD
Kyiv, Ukraine

Nur A. Hatab, DMD, PhD
Ras Al Khaimah, United Arab Emirates

Vladyslav P. Iefymenko, PhD
Kyiv, Ukraine

Michael T. Kase, DMD
Birmingham, Alabama, United States

Andrey V. Kopchak, ScD
Kyiv, Ukraine

Olindo Massarelli, MD, PhD, FEBOMFS
Sassari, Italy

Anastasiya Quimby, DDS, MD
Fort Lauderdale, Florida, United States

Daniel Robles Cantero, DDS, MSc, PhD
Madrid, Spain

Nataliia O. Savychuk, ScD
Kyiv, Ukraine

Peter Stanko, MD, PhD
Bratislava, Slovak Republic

Rostyslav M. Stupnytskyi, ScD
Kyiv, Ukraine

Olexander O. Tymofieiev, ScD
Kyiv, Ukraine

Natalia O. Ushko, ScD
Kyiv, Ukraine

Andrew Yampolsky, DDS, MD
Philadelphia, Pennsylvania, United States

Web & Social Media Editor

João L. Monteiro, DDS, PhD
Boston, Massachusetts, United States

Resident Ambassador

Marcus Hwang, DDS, MD
Portland, Oregon, United States

Managing Editor

Ievgen I. Fesenko, PhD
Kyiv, Ukraine



TANTUM VERDE®

INFORMATION LEAFLET
for the medicinal product

Composition:

active substance: benzydamine hydrochloride;

100 mL of solution contain benzydamine hydrochloride 0.15 g;

excipients: ethanol 96%, glycerol, methyl parahydroxybenzoate (E 218), flavor (menthol), saccharin, sodium hydrocarbonate, Polysorbate 20, Quinoline Yellow (E 104), Patent Blue V (E 131), purified water.

Dosage form. Oromucosal solution.

Basic physical and chemical properties: a clear green liquid with a typical mint flavor.

Pharmacotherapeutic group. Dental preparations. Other agents for local oral treatment.

ATC code: A01A D02.

Pharmacological properties.

Pharmacodynamics.

Benzydamine is a non-steroidal anti-inflammatory drug (NSAID) with analgesic and antiexudative properties.

Clinical studies have shown that benzydamine is effective in the relief of symptoms accompanying localized irritation conditions of the oral cavity and pharynx. Moreover, benzydamine has anti-inflammatory and local analgesic properties, and also exerts a local anesthetic effect on the oral mucosa.

Pharmacokinetics.

Absorption through the oral and pharyngeal mucosa has been proven by the presence of measurable quantities of benzydamine in human plasma. However, they are insufficient to produce any systemic pharmacological effect. The excretion occurs mainly in urine, mostly as inactive metabolites or conjugated compounds.

When applied locally, benzydamine has been shown to cumulate in inflamed tissues in an effective concentration

due to its ability to permeate through the mucous membrane.

Clinical particulars.

Indications.

Symptomatic treatment of oropharyngeal irritation and inflammation; to relieve pain caused by gingivitis, stomatitis, pharyngitis; in dentistry after tooth extraction or as a preventive measure.

Contraindications.

Hypersensitivity to the active substance or to any other ingredients of the product.

Interaction with other medicinal products and other types of interaction.

No drug interaction studies have been performed.

Warnings and precautions.

If sensitivity develops with long-term use, the treatment should be discontinued and a doctor should be consulted to get appropriate treatment.

In some patients, buccal/pharyngeal ulceration may be caused by severe pathological processes. Therefore, the patients, whose symptoms worsen or do not improve within 3 days or who appear feverish or develop other symptoms, should seek advice of a physician or a dentist, as appropriate.

Benzydamine is not recommended for use in patients hypersensitive to acetylsalicylic acid or other non-steroidal anti-inflammatory drugs (NSAIDs).

The product can trigger bronchospasm in patients suffering from or with a history of asthma. Such patients should be warned of this.

For athletes: the use of medicinal products containing ethyl alcohol might result in positive antidoping tests considering the limits established by some sports federations.

Use during pregnancy or breast-feeding

No adequate data are currently available on the use of benzydamine in pregnant and breastfeeding women. Excretion of the product into breast milk has not been studied. The findings of animal studies are insufficient to make any conclusions about the effects of this product during pregnancy and lactation.

The potential risk for humans is unknown.

TANTUM VERDE should not be used during pregnancy or breast-feeding.

Effects on reaction time when driving or using machines

When used in recommended doses, the product does not produce any effect on the ability to drive and operate machinery.

Method of administration and doses.

Pour 15 mL of TANTUM VERDE solution from the bottle into the measuring cup and gargle with undiluted or diluted product (15 mL of the measured solution can be diluted with 15 mL of water). Gargle 2 or 3 times daily. Do not exceed the recommended dose.

Children.

The product should not be used in children under 12 years due to a possibility of ingestion of the solution when gargling.

Overdosage.

No overdose has been reported with benzydamine when used locally. However, it is known that benzydamine, when ingested in high doses (hundreds times higher than those possible with this dosage form), especially in children, can cause agitation, convulsions, tremor, nausea, increased sweating, ataxia, and vomiting. Such acute overdose requires immediate gastric lavage, treatment of fluid/salt imbalance, symptomatic treatment, and adequate hydration.

Adverse reactions.

Within each frequency group, the undesirable effects are presented in order of their decreasing seriousness.

Adverse reactions are classified according to their frequency: very common ($\geq 1/10$); common ($\geq 1/100$ to $<1/10$); uncommon ($\geq 1/1,000$ to $<1/100$); rare ($\geq 1/10,000$ to $<1/1,000$); very rare ($<1/10,000$); frequency unknown (cannot be estimated from the available data).

Gastrointestinal disorders: rare – burning mouth, dry mouth; *unknown* – oral hypesthesia, nausea, vomiting, tongue edema and discoloration, dysgeusia.

Immune system disorders: rare – hypersensitivity reaction, *unknown* – anaphylactic reaction.

Respiratory, thoracic and mediastinal disorders: very rare – laryngospasm; *unknown* – bronchospasm.

Skin and subcutaneous tissue disorders: uncommon – photosensitivity; very rare – angioedema; *unknown* – rash, pruritus, urticaria.

Nervous system disorders: *unknown* – dizziness, headache.

TANTUM VERDE contains methyl parahydroxybenzoate, which can cause allergic reactions (including delayed-type reactions).

Shelf life. 4 years.

Storage conditions.

Do not store above 25°C. Keep out of reach of children.

Packaging.

120 mL of solution in a bottle with a measuring cup; 1 bottle per cardboard box.

Dispensing category.

Over-the-counter medicinal product.

Manufacturer.

Aziende Chimiche Riunite Angelini Francesco A.C.R.A.F. S.p.A., Italy.

Location of the manufacturer and its business address.
Via Vecchia del Pinocchio, 22 – 60100 Ancona (AN), Italy.

Date of the last revision of the text.

September 26, 2018.

Information leaflet is

APPROVED by

Order of the

Ministry of Health of Ukraine

No. 636 dated 01.10.2015

Registration Certificate

No. UA/3920/01/01

State Registration

JULY 2025 • VOLUME 9 • ISSUE 7
www.djournal.org


Міністерство юстиції України
СВІДОЦТВО
про державну реєстрацію
друкованого засобу масової інформації

Серія KB № 24951-14891 ПП

«Журнал діагностики та лікування оральної і щелепно-лицевої патології»
(назва видання державною мовою)
«Journal of Diagnostics and Treatment of Oral and Maxillofacial Pathology»
(назва видання іншою мовою (мовами))

Вид видання журнал
(газета, журнал, бюлетень, збірник, альманах, календар, дайджест)

Статус видання вітчизняне
(вітчизняне, спільне)

Мова (мови) видання змішаними мовами: українська, англійська


Вид видання за цільовим призначенням наукове, науково-виробниче
(громадсько-політичне, наукове, навчальне, інформаційне, рекламне (понад 40 відсотків обсягу одного номера – реклама), еротичне тощо)

Обсяг, періодичність 23,1 ум. друк. арк., формат А4 (210x297мм), 1 раз на місяць

Сфера розповсюдження та категорія читачів загальнодержавна, зарубіжна
лікарі-стоматологи-хірурги; лікарі ультразвукової діагностики, лікарі-рентгенологи, лікарі-патологоанатоми, студенти, лікарі-інтерни, слухачі, аспіранти, докторанти, наукові, науково-педагогічні та педагогічні працівники закладів вищої освіти та наукових установ

Засновник (співзасновники) Національний університет охорони здоров'я України імені П.Л. Шупика (код за ЄДРПОУ 01896702), Приватний вищий навчальний заклад «Київський медичний університет» (код за ЄДРПОУ 16478809), Товариство з обмеженою відповідальністю «ОМФ ПАБЛІШІНГ» (код за ЄДРПОУ 40493077)

Програмні цілі (основні принципи) журнал присвячений сучасним проблемам діагностики і лікування в хірургічній стоматології та щелепно-лицевій або тематична спрямованість хірургії

Перший заступник Міністра  **Євгеній ГОРОВЕЦЬ**

10.08.2021
(дата реєстрації)



FIGURE 1. Certificate of State Re-Registration of the Print Mass Media (journal) in the Ministry of Justice of Ukraine as of 2021. The *Journal* was registered for the first time in 2016 under the title *Diagnostics and Treatment of Oral and Maxillofacial Pathology*. The next re-registration took place in 2019.

State Registration

JULY 2025 • VOLUME 9 • ISSUE 7
www.dtjournal.org


УКРАЇНА

**НАЦІОНАЛЬНА РАДА УКРАЇНИ
З ПИТАНЬ ТЕЛЕБАЧЕННЯ І РАДІОМОВЛЕННЯ**

01601, Київ, вул. Прорізна, 2, тел. (38044) 278-68-32, факс (38044) 278-74-90
<http://www.nrada.gov.ua>, E-mail: nrada@nrada.gov.ua

_____ 20 ____ р. № _____
на № _____

ВИТЯГ*
З РЕЄСТРУ СУБ'ЄКТІВ У СФЕРІ МЕДІА - РЕЄСТРАНТІВ

Виданий ТОВАРИСТВУ З ОБМЕЖЕНОЮ ВІДПОВІДАЛЬНІСТЮ «ОМФ ПАБЛІШИНГ», м. Київ, код ЄДРПОУ 40493077
(найменування суб'єкта у сфері медіа, код згідно з ЄДРПОУ)

Ідентифікатор медіа R30-04318

Логотип, позивні або назва медіа «Журнал діагностики та лікування оральної і щелепно-лицевої патології» / «Journal of Diagnostics and Treatment of Oral and Maxillofacial Pathology»

Суб'єкт у сфері друкованих медіа, вид друкованого медіа –журнал
(суб'єкт, що здійснює мовлення без використання радіочастотного спектра (лінійне медіа, вид та технологія мовлення), суб'єкт у сфері аудіовізуальних медіа на замовлення (нелінійний медіа-сервіс, технологія, що застосовується для надання сервісу), провайдер аудіовізуальних сервісів (технологія, що застосовується для надання доступу), провайдер платформ спільного доступу до відео (технологія, що застосовується для надання сервісу), суб'єкт у сфері друкованих медіа (вид друкованого медіа), суб'єкт у сфері онлайн-медіа (вид онлайн медіа))

Рішення від 11.04.2024 № 1225
(дата і номер рішення Національної ради про реєстрацію; дата і номер рішення про внесення останніх змін до Реєстру)

М. П.
Перший заступник голови
Валентин КОВАЛЬ

Відповідальний секретар
Олена НІЦКО



* до введення в дію Електронного кабінету (за вимогою суб'єкта)

A

FIGURE 2. The *Journal* is registered by the National Council of Television and Radio Broadcasting of Ukraine as print media **(A)** (Decision of April 11, 2024, No. 1225) and online media **(B)** (Decision of April 25, 2024, No. 1454). The identifier of print media is R30-04318 and the identifier of online media is R40-04708. **(Fig 2 continued on next page.)**

State Registration

JULY 2025 • VOLUME 9 • ISSUE 7
www.dtjournal.org


УКРАЇНА

**НАЦІОНАЛЬНА РАДА УКРАЇНИ
З ПИТАНЬ ТЕЛЕБАЧЕННЯ І РАДІОМОВЛЕННЯ**
01601, Київ, вул. Прорізна, 2, тел. (38044) 278-68-32, факс (38044) 278-74-90
<http://www.nrada.gov.ua>, E-mail: nrada@nrada.gov.ua

_____ 20 __ р. № _____
на № _____

ВИТЯГ*
З РЕЄСТРУ СУБ'ЄКТІВ У СФЕРІ МЕДІА - РЕЄСТРАНТІВ

Виданий ТОВАРИСТВУ З ОБМЕЖЕНОЮ ВІДПОВІДАЛЬНІСТЮ «ОМФ ПАБЛІШИНГ», м. Київ, код ЄДРПОУ 40493077
(найменування суб'єкта у сфері медіа, код згідно з ЄДРПОУ)

Ідентифікатор медіа R40-04708

Логотип, позивні або назва медіа «Journal of Diagnostics and Treatment of Oral and Maxillofacial Pathology» / «Журнал діагностики та лікування оральної і щелепно-лицевої патології»

Суб'єкт у сфері онлайн-медіа, вид онлайн-медіа – журнал
(суб'єкт, що здійснює мовлення без використання радіочастотного спектра (лінійне медіа, вид та технологія мовлення), суб'єкт у сфері аудіовізуальних медіа на замовлення (нелінійний медіа-сервіс, технологія, що застосовується для надання сервісу), провайдер аудіовізуальних сервісів (технологія, що застосовується для надання доступу), провайдер платформ спільного доступу до відео (технологія, що застосовується для надання сервісу), суб'єкт у сфері друкованих медіа (вид друкованого медіа), суб'єкт у сфері онлайн-медіа (вид онлайн медіа))

Рішення від 25.04.2024 № 1454
(дата і номер рішення Національної ради про реєстрацію; дата і номер рішення про внесення останніх змін до Реєстру)

М. П. _____

Перший заступник голови  **Валентин КОВАЛЬ**

Відповідальний секретар **Олена НІЦКО**

* до введення в дію Електронного кабінету (за вимогою суб'єкта)

B

FIGURE 2 (continued). The *Journal* is registered by the National Council of Television and Radio Broadcasting of Ukraine as print media (**A**) (Decision of April 11, 2024, No. 1225) and online media (**B**) (Decision of April 25, 2024, No. 1454). The identifier of print media is R30-04318 and the identifier of online media is R40-04708. (**Fig 2 continued on next page.**)

Contents

of the Volume 9 • Issue 7 • July 2025

Article Type	Article Number	Pages	Description
		A1	Publisher & Editorial Office Information
		A2	Editorial Board
		A5-A7	State Registration
		A8	Contents, Courtesy, & Erratum
CASE REPORT/ TECHNIQUE	100307	1-11	A Case Report of Oral Mucosa Metastasis of Renal Clear Cell Carcinoma Mimicking Benign Neoplasm: Possibilities and Limitations of the Ultrasonography Olha S. Cherniaka, Valentyn H. Demidov, Oleksandra Petrychenko, & Viacheslav P. Blyzniuk



COURTESY

The *Journal*'s cover images are courtesy Olindo Massarelli, MD, PhD, FEBOMFS.

The images were taken from this article: Massarelli O, Meloni SM. The “beveled one-and-a-half-barrel” fibula transplant with virtual surgical planning and CT-guided implant surgery for prosthetic rehabilitation in posterior mandible defects: a pictorial essay. *J Diagn Treat Oral Maxillofac Pathol* **2022**;6(3):39–59.

<https://doi.org/10.23999/j.dtopm.2022.3.3>



CASE REPORT/TECHNIQUE

USG

A Case Report of Oral Mucosa Metastasis of Renal Clear Cell Carcinoma Mimicking Benign Neoplasm: Possibilities and Limitations of the Ultrasonography

Olha S. Cherniak^{a,*}, Valentyn H. Demidov^b, Oleksandra Petrychenko^c, & Viacheslav P. Blyzniuk^d

SUMMARY

Oral metastases are rare, accounting for approximately 1-1.5% of all oral malignancies. Therefore, it is important for clinicians to create diagnostic data for early verification of this formidable pathological process. The most common primary tumors that metastasise to the oral cavity are: lung (21.1%), liver (12.3%), breast (10.5%), kidney (10.5%) and colorectal (8.8%). We hope that the publication of case described below will complement the international literature and bring ultrasound diagnostics to a new level. A 49-year-old Caucasian female with a growing neoplasm in the oral cavity and its mushroom-shaped attachment to the mucosa of the mandibular alveolar process presented for surgical examination. The neoplasm clinically mimicked a benign pathology. The anterior part of the neoplasm was examined using ultrasonography with the presentation of grayscale and color Doppler sonograms. It is worth noting that the vestibular (anterior, smaller) part of the metastasis in this case did not have a characteristic sign for metastases in other parts of the body, namely, there was no vascularization on color Doppler ultrasonography. The article also provides clinical and orthopantomography data. The limitations of the linear probe for a complete ultrasound assessment of the neoplasms located on the lingual or possibly palatal sides of the alveolar processes were analyzed. A comparative analysis of a linear and a hockey-stick probes is presented. After conducting our analysis of the literature, we can state that this article presents for the first time data on ultrasound examination of oral mucosa metastasis of renal clear cell carcinoma. Further investigation of each oral metastasis using intra-

^a Doctor of Ultrasound Diagnostics; Head, Department of Ultrasound, Regional Diagnostic Center, Polyclinic, Kyiv Regional Clinical Hospital, Public Non-profit Enterprise of the Kyiv Region Council, Kyiv, Ukraine.

^b Doctor-Stomatologist-Surgeon, Center of Maxillofacial Surgery and Dentistry, Kyiv Regional Clinical Hospital, Public Non-profit Enterprise of the Kyiv Region Council (place of work at moment of material collection), Kyiv, Ukraine.

^c Doctor-Stomatologist-Surgeon, Center of Maxillofacial Surgery and Dentistry, Kyiv Regional Clinical Hospital, Public Non-profit Enterprise of the Kyiv Region Council (place of work at moment of material collection), Kyiv, Ukraine; Owner, Unique Aesthetic, Batumi, Georgia.

^d Independent Researcher; Doctor-Stomatologist-Surgeon, Center of Maxillofacial Surgery and Dentistry, Kyiv Regional Clinical Hospital, Public Non-profit Enterprise of the Kyiv Region Council (place of work at moment of material collection), Kyiv, Ukraine.

* Corresponding author's address: Office 418, Polyclinic, Kyiv Regional Clinical Hospital, 1 Zahorivska Street, Kyiv 04107, Ukraine.
E-mail: cherniak.os@gmail.com (Olha Cherniak).
ORCID: <https://orcid.org/0000-0002-6372-0043>.

Article type: Case report/technique.

Please cite this article as: Cherniak OS, Demidov VH, Petrychenko O, Blyzniuk VP. A case report of oral mucosa metastasis of renal clear cell carcinoma mimicking benign neoplasm: Possibilities and limitations of the ultrasonography. J Diagn Treat Oral Maxillofac Pathol. 2025 Jul;9(7):100307.

At the time of publication of this article, the journal's impact factor (Exaly) is 0.2. Each author of this article is awarded 20 CPD points according to the Order of the Ministry of Healthcare of Ukraine of February 22, 2019, No. 446. The bibliographic reference is a supporting document.

The acronym "USG" in the upper right icon means that this article describes images of ultrasonography (USG).

Manuscript received 03 June 2025

Accepted 26 June 2025

Available online 31 July 2025

<https://doi.org/10.23999/j.dtopm.2025.7.100307>

© 2025 The Author(s). Published by OMF Publishing LLC.

Website: <https://omfpublishing.com>.

This is an open access article under the CC BY-NC-SA license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>).

oral direct ultrasound examination is extremely important for systematizing the ultrasound variants of this pathology. In turn, this will allow clinicians to more likely establish a correct preliminary diagnosis before biopsy and treat this pathology in specialized departments.

KEY WORDS

Oral metastasis, renal clear cell carcinoma, ultrasonography, linear probe, linear hockey-stick probe

Takeaways

Question: Is there a solution for ultrasound imaging and diagnosis of neoplasms located on the lingual and palatal side of the alveolar processes?

Findings: Considering that the use of direct ultrasonography from the lingual and palatal sides of the alveolar ridge is difficult to access with a linear probe, we suggest using a linear hockey stick probe. This is especially important in the early diagnosis of oncological processes.

Meaning: Providing the possibility of direct contact of the hockey-stick linear probe with pathological tissues will allow for their full ultrasound examination while preserving grayscale, color, and power Doppler sonograms.

INTRODUCTION

Metastatic tumors to the oro-facial region are uncommon and account for approximately 1-1.5% of all malignant oral tumors [1, 2]. The most common primary tumors that metastasise to the oral cavity are: lung (21.1%), liver (12.3%), breast (10.5%), kidney (10.5%) and colorectal (8.8%) [3]. That is, out of 5 different groups of organs from which cancer can metastasize to the oral cavity, kidney cancer ranks 4th in frequency. Renal cell carcinoma (RCC) is the most common type of primary renal cancer, and clear-cell is the most common subtype [4]. The ultrasonography (USG) patterns of malignant lymph nodes are well described [5]. Although there are quite a few published cases of renal cell carcinoma metastasis to the oral cavity [2, 6-15], we were unable to find a single case with documented USG of such metastases. This may be due to the rarity of the pathology itself, the absence of USG in the examination protocol, and the lack of a special ultrasound probe for insertion into the oral cavity.

So, the purpose of this report is to highlight the case of oral mucosa metastasis of renal clear cell carcinoma, present the results of first to our knowledge ever reported ultrasound examination of such pathologic condition.

CASE REPORT

In May 19, 2015, a 49-year-old Caucasian female

was referred to the Center of Maxillofacial Surgery and Dentistry with complaints of the presence of a neoplasm on the mucosa of the alveolar process of the mandible. According to the patient, this neoplasm has existed for the past few months and is growing. When taking the history, the patient did not mention any surgeries or treatments for RCC.

Clinical examination revealed the presence of a neoplasm on the lingual side of the alveolar process of the mandible that had grown vestibularly between the left central and lateral incisors (Fig 1). The neoplasm was mushroom-shaped, pale pink in color, elastic upon palpation, and had a pedicle. The mucosa adjacent to the neoplasm is unchanged in color and structure. The neoplasm measured 1.8 x 1.1 x 1.2 cm at the lingual aspect of the lower teeth and 0.34 x 0.38 cm at the vestibular aspect.

Orthopantomography (Fig 2) showed a minor area of alveolar ridge resorption in the area between the lower left lateral incisor and canine.

An extraoral ultrasound examination was performed by O.S.C., an expert physician (>18 years of experience in head and neck USG), on the day of admission in the department. A HD11 XE USG system (2007, the Philips, Amsterdam, the Netherlands) equipped with a linear L12-3 transducer (frequency range: 3.0-12.0 MHz; aperture: 38 mm) was used for the examination, fixing a 3-cm of ultrasonography (penetration) depth.

Anterior part of the neoplasm at the vestibular

aspect of the lower teeth measured 0.34 x 0.38 cm (Fig 3). Upon the gray-scale USG the anterior part portion of the neoplasm vizualized as homogenous hypoechoic lesion with no signs of calcifications. Color Doppler USG showed no vascularity within the vestibular portion of the neoplasm (Fig 4).

Unfortunately, this ultrasound machine in our hospital was equipped only with a linear and convex probe but was not equipped with a hockey-stick probe. Application of linear probe “hockey stick” is a solution for ultrasound imaging and diagnostics of neoplasms located on the lingual and palatal sides of

the teeth, i.e., for the intraoral USG.

This ultrasound machine that we used to examine this patient, namely HD11 XE, is suitable for a transducer such as L15-7io hockey stick ultrasound probe (the Philips, Amsterdam, the Netherlands; frequency range: 7.0-15.0 MHz; aperture: 23 mm).

An excisional biopsy of the neoplasm was performed under general anesthesia. The histopathological diagnosis of metastasis of renal clear cell carcinoma was made by experienced pathologists. The patient was referred for further treatment to an oncologist.




FIGURE 1. A neoplasm attached to the mandibular mucosa arising between the left central and lateral incisors. The neoplasm is mushroom-shaped and has two lobes, an anterior lesser (*white arrowhead*) and a posterior greater (*black arrow*).




FIGURE 2. Preoperative orthopantomography.




FIGURE 3. Preoperative gray-scale ultrasonography. Vestibular (i.e., anterior) portion of the metastasis is indicated by "+" and "x" calipers. 3.1, lower left central incisor; 3.2, lower left lateral incisor. Penetration depth: 3.0 cm.




FIGURE 4. Preoperative color Doppler ultrasonography. Vestibular (i.e., anterior) portion of the metastasis is indicated by *open arrowhead*. 3.1, lower left central incisor; 3.2, lower left lateral incisor. Penetration depth: 3.0 cm.

DISCUSSION AND TECHNIQUE

Metastasis is a spread of cancer cells from the place where they first formed to another part of the body [16]. In metastasis, cancer cells break away from the original (primary) tumor, travel through the blood or lymph system, and form a new tumor in other organs or tissues of the body [16].

In metastatic nodes, upon the gray-scale USG assessment of size, shape, borders, echogenicity, echogenicity of the hilum, presence of intranodal necrosis, calcifications, and ancillary features (matting of the nodes and adjacent soft tissue oedema) is performed [5]. Upon the Doppler USG of the metastatic nodes vascular distribution and resistance are examining [5].

USG of skin metastasis (arms or buttock localization) of carcinomas is already presented in the literature [17-19]. Key characteristics according to Corominas et al (2015) were next: (1) metastasis of the prostate adenocarcinoma was iso/hypoechoic on gray scale sonograms, moderately well-defined, and was joined to a vessel that fed it, and (2) the marked vascularity on color/power Doppler [17].

USG pattern of metastasis according to Martínez-Morán et al (2018) was next: A heterogeneous solid-cystic lesion in the dermis and extending into the subcutaneous tissue [18]. The lesion also had anechoic central areas admixed with hyperechoic areas with well-defined borders [18]. It displayed posterior reinforcement and power Doppler USG showed peripheral vascular poles [18].

Bai et al (2022) reported the cutaneous metastasis of RCC [19]. Superficial USG of the right arm in their case showed (1) an oval, hypo-echoic nodule with well-defined margins and no shadowing in the subcutaneous fat, (2) extensive vascularity [19].

Pickhardt and Pickard (2003) reported the delayed thyroid metastasis from RCC with jugular vein extension and color Doppler showed prominent metastatic tumor vascularity [20].

Even though many sources describe the ultrasound picture of carcinoma metastases, USG patterns of the oral metastasis of the RCC are not described. In any case, we were unable to describe a single case with an ultrasound picture of oral metastasis of the RCC. It is important to know that *renal clear cell carcinoma* is also terms as *hypernephroma*, *Grawitz tumor*, and *renal cell carcinoma* (Makos and Psomadakis [2009]) [7]. Metastasis of the renal clear cell carcinoma spread

to different organs including bones, the brain, the pancreas, the gallbladder, lymph nodes [21], and oral cavity [2, 6-15].

In the case of benign neoplasm/primary malignant tumor/metastatic tumor in the oral cavity, clinicians can visually assess their size, shape, surface, and structure with and without palpation. In addition to the clinical examination, USG will allow to assess the internal structure, the boundaries of the neoplasm, the depth of invasion into adjacent tissues and vascularization. Some cases of USG in the intraoral benign and malignant tumors are well described by Orhan and Ünsal (2021) [22].

Again, we reviewed a significant number of sources describing oral metastases of RCC and found no ultrasound images. In particular, van der Waal et al (2003) in their report on 24 cases of oral metastases indicate that in four cases the site of the primary tumor was the kidney, and the site of oral metastases was the maxilla, mandible, soft palate, and buccal mucosa [2]. Histological type in these 4 cases was clear cell carcinoma [2]. The age of these patients ranged from 48 to 67 years, and the sex ratio was 50/50% [2].

Hirshberg et al (2008) performed important analysis of 673 intraoral metastasis of carcinomas from different primary sites [23]. The jawbones (n = 36) were more frequently affected by metastasis of RCC than the oral soft tissues (n = 29). For example, Kelles et al (2012) presented a 59-year-old woman who had left nephrectomy because of renal cell carcinoma three years ago referred with trismus and a metastasis on her temporomandibular joint [24].

We consider the following cases noteworthy, as they allow for clinical comparison with our case. Sastre et al (2005) described a 64-year-old male with a metastasis along the right alveolar ridge of the mandible [6].

Makos and Psomadakis (2009) published a case of a 63-year-old male with a maxillary epulis-like metastasis from the right central maxillary incisor to the first right maxillary premolar [7].

Dashow et al (2011) highlighted a case of 67-year-old gentleman metastatic clear cell renal carcinoma of the palate mimicking noninvoluting congenital hemangioma [8]. Metastasis visualized as a nonhealing, exophytic lesion on his soft-hard palate, which the patient had been aware of for approximately one year. The patient had a history of a renal cell carcinoma diagnosis in 1991, which had been treated with radical right nephrectomy.

Milner et al (2014) reported a 67-year-old male with a metastasis on the hard palate mucosa which looked somewhat like a case of Dashow et al (2011) [8, 9].

Selvi et al (2016) presented three synchronous atypical metastases of RCC to the maxillary gingiva, scalp and the distal phalanx of the fifth digit in a 51-year-old male [10].

Vasilyeva et al (2018) described a metastasis of RCC as a tan-red exophytic, lobulated mass of the maxillary anterior facial gingiva with a separate, similar appearing smaller lesion identified in the right maxillary vestibule [25].

Nisi et al (2020) reported two males, 61- and 71-year-old, with oral metastases located on the left tongue and buccal mucosa [11].

Kovalski et al (2020) detailed double lobe nodule covering the gingiva and alveolar ridge on the left side of the maxilla in a 63-year-old male [12]. The metastasis measured 5 x 3 cm, had a reddish colouration, was covered by a purulent membrane, bled easily when touched and had a foul odour.

Patel et al (2020) delineate oral metastasis in a 59-year-old female as a pink-red, oval, ulcerated lesion with a white pseudomembranous surface measuring approximately 3.8 x 2.5 x 1.7 cm attached to the left buccal mucosa via a pedunculated stalk [14].

Intraoral metastasis of RCC is also described as exophytic tumor mass, sized about 6 x 3 cm. In the anteroposterior direction, it extended from the mandible level, involving the alveolar portion, to the retromolar triangle on the right; craniocaudally, the lesion extended from the lower fornix, over the top of the alveolar process, crossing it over, and descending toward the floor of the mouth (Stojanovic et al, 2020). In the described case in a 53-year-old male was also noted a enlargement of the right half of the face [13].

Khushbu et al (2024) reported metastasis to the upper gingiva in a 52-year-old female [26].

Metastasis to the upper maxilla in a 79-year-old male from the study by Nakako et al (2025) was clinically similar to our case [15].

Interestingly, dermatologic USG requires using multifrequency linear or compact linear (hockey-stick) probes [27]. Probes working with their upper-frequency range between 15 and 24 MHz allow displaying the skin layers and their abnormalities with sufficiently high-quality images [27]. Ultrasound in skin cancer and skin metastases

are illustrated in the study by Wortsman (2024) [28].

Healthcare professionals may attempt to examine various areas of the oral cavity both extra- and intraoral USG.

We consider it advisable and indication to use a hockey stick probe for intraoral ultrasound scanning of all intraoral lesions. A comparison of the characteristics of linear and hockey-stick probes from one manufacturer, namely GE (GE Healthcare Japan Corporation, Tokyo, Japan), is presented in Table 1.

The use of “hockey stick” probe is also very useful for young and small patients due to the lack of space for the use of the regular linear probe (Guimarães Ferreira Fonseca et al, 2024) [29]. Synonymous name for the linear probe “hockey stick” is a *small-footprint linear array transducer*.

The use of a linear transducer for intraoral ultrasound scanning and diagnosis of pathology that is in close proximity to the transducer surface is possible. For example, this is possible for diagnosing tongue pathology, provided there is sufficient mouth opening and/or some tongue mobility [30-33]. However, the use of a hockey-type transducer is the most rational, in particular for cases like ours.

The lower level of floor of mouth and the lower margin of the mandible can be assessed perfectly with extraoral ultrasound using a linear transducer [34-36]. Ultrasonographic capabilities in examining the floor of the mouth and the root of the tongue using linear and curved probes are presented in the study by Abbott et al (2020) [32]. The clinicians should understand that penetration depth is the highest distance between the probe and the tissues which can still be visualized without image noise [37].

Thus, our proposal is to use a hockey-stick probe [38, 39] in cases with a pathology location like our case. As an option, we suggest using the following transducer: Linear probe “hockey stick” GE L8-18i-RS (GE Healthcare Japan Corporation, Tokyo, Japan).

One of the important points in obtaining high-quality informative sonograms when using a linear probe is the proximity of the target diagnostic area (area with a pathological focus) to the surface of the probe [40]. Since with a remote localization of the pathology area, the diagnostic ability decreases due to the likelihood of the presence of anatomical objects/foreign bodies that can produce artifacts and due to the technical limitations of the ultrasound

TABLE 1. Comparison of Ultrasound Probes (GE Healthcare Japan Corporation, Tokyo, Japan) for Direct and Indirect Intraoral Ultrasonography in Adults and Children.

	Features	Linear probe	Linear probe “hockey stick”
1	Probe view		
2	Name	GE ML6-15-RS Linear Array transducer	GE L8-18i-RS Linear Array transducer
3	Footprint, cm	Flat and longer: 6.07 cm x 1.6 cm	Flat and shorter: 3.48 cm x 1.11 cm
4	Frequency range	4.0-15.0 MHz	6.7-18.0 MHz
5	Compatibility	GE Voluson S8, S10; GE Logiq P9; GE Vivid iq	GE Logiq P9; GE Logiq e; GE Vivid iq; GE Venue Go
6	Possible direct approach (i.e., intraoral approach) to the oral area of examination	Anterior and middle portion of the tongue in case it is possible to protrude it and open the mouth; vestibular aspect of the alveolar processes of the jaws	Anterior, middle, posterior, and lateral portions of the tongue in case it is possible to protrude it and open the mouth; palate; palatal, lingual, and vestibular aspects of the alveolar processes of the jaws
7	Possible distant approach (i.e., extraoral approach) to the oral area of examination (transcervical, facial, perioral/translabial)	Floor of the mouth	Floor of the mouth
8	Depth of ultrasonography (depth of field), cm	8 cm	10 cm

probe itself. That is, ideally, the area with pathology should be located at a depth of 0 to 3.0-6.0 cm relative to the footstep of the transducer.

According to the literature (Cheng et al, 2000; Dashow et al, 2011), the 5-year survival rate for RCC after nephrectomy is 60% to 75%; with a solitary oral metastasis, the treatment is radical excision and nephrectomy, with a 5-year survival rate of 35% [41, 8]. The prognosis for patients with multiple RCC metastases is poor, with a 5-year survival rate of 0% to 7%.

In summary, metastasis to the oral cavity is rare, accounting for approximately 1-1.5% of all oral malignancies [1, 42, 43]. Therefore, it is extremely important to develop diagnostic data for the verification of such metastases, including ultrasound data. It is worth noting that the vestibular (anterior,

smaller) part of the metastasis in this case did not have a characteristic sign for metastases in other parts of the body, namely, there was no vascularization on color Doppler ultrasonography. Hypothetically, this can be explained either by the initial stage of metastasis development [44] and the location in relation to soft tissues and bony structures.

CONCLUSION

Further investigation of each oral metastasis using intraoral direct ultrasound examination is extremely important for systematizing the ultrasound variants of this pathology. In turn, this will allow clinicians to more likely establish a correct preliminary diagnosis before biopsy and treat this pathology in specialized departments.

DISCLOSURE

The authors have no financial interest to declare in relation to the content of this article.


PATIENT CONSENT

Informed consent was obtained from the patient at the time of presentation.

REFERENCES (44)

- Kumar G, Manjunatha B. Metastatic tumors to the jaws and oral cavity. *J Oral Maxillofac Pathol.* **2013** Jan;17(1):71-5.
<https://doi.org/10.4103/0973-029X.110737>
- van der Waal RI, Buter J, van der Waal I. Oral metastases: report of 24 cases. *Br J Oral Maxillofac Surg.* **2003** Feb;41(1):3-6.
[https://doi.org/10.1016/s0266-4356\(02\)00301-7](https://doi.org/10.1016/s0266-4356(02)00301-7)
- Liu Y, Vargo RJ, Bilodeau EA. Analytic survey of 57 cases of oral metastases. *J Oral Pathol Med.* **2018** Mar;47(3):275-280.
<https://doi.org/10.1111/jop.12672>
- Silver E, Roudakova K, Bial N, Daniel D. Cutaneous metastasis of renal cell carcinoma to the cheek: A case report and literature review. *Am J Case Rep.* **2021** Mar 26;22:e928999.
<https://doi.org/10.12659/AJCR.928999>
- Ahuja AT, Ying M, Ho SY, et al. Ultrasound of malignant cervical lymph nodes. *Cancer Imaging.* **2008** Mar 25;8(1):48-56.
<https://doi.org/10.1102/1470-7330.2008.0006>
- Sastre J, Naval L, Muñoz M, Gamallo C, Diaz FJ. Metastatic renal cell carcinoma to the mandible. *Otolaryngol Head Neck Surg.* **2005**;132(4):663-664.
- Makos CP, Psomaderis K. A literature review in renal carcinoma metastasis to the oral mucosa and a new report of an epulis-like metastasis. *J Oral Maxillofac Surg.* **2009**;67(3):653-660.
<https://doi.org/10.1016/j.joms.2008.10.006>
- Dashow JE, Gemmete JJ, McHugh JB, Helman JL. Metastatic clear cell renal carcinoma of the palate mimicking noninvoluting congenital hemangioma. *J Oral Maxillofac Surg.* **2011**;69(6):1836-1841.
<https://doi.org/10.1016/j.joms.2010.07.050>
- Milner P, Janas A, Grzesiak-Janias G. Clear cell renal carcinoma metastasis in the oral cavity – Case report. *J Pre Clin Clin Res.* **2014** Jun;8(2):127-129.
<https://doi.org/10.26444/jpccr/71484>
- Selvi F, Faquin WC, Michaelson MD, August M. Three synchronous atypical metastases of clear cell renal carcinoma to the maxillary gingiva, scalp and the distal phalanx of the fifth digit: A case report. *J Oral Maxillofac Surg.* **2016** Jun;74(6):1286.e1-1286.e12869.
<https://doi.org/10.1016/j.joms.2016.01.054>
- Nisi M, Izzetti R, Graziani F, Gabriele M. Renal cell carcinoma metastases to the oral cavity: Report of 2 cases and review of literature. *J Oral Maxillofac Surg.* **2020** Sep;78(9):1557-1571.
<https://doi.org/10.1016/j.joms.2020.04.001>
- Kovalski LNS, Ribeiro JT, Martins MD, et al. A rare case of oral metastasis of renal clear cell carcinoma: case report and review of literature. *J Oral Diagn.* **2020** Jan;5:1-9.
<https://doi.org/10.5935/2525-5711.20200006>
- Stojanovic M, Krasic D, Trajkovic M, Petrovic V. Rare renal cell carcinoma metastasis to mandibular gingiva: A case report and literature review. *Niger J Clin Pract.* **2020**;23(10):1483-1486.
https://doi.org/10.4103/njcp.njcp_55_19
- Patel S, Barros J, Nwizu NN, Ogbureke KUE. Metastatic renal cell carcinoma to the oral cavity as first sign of disease: A case report. *Clin Case Rep.* **2020**;8(8):1517-1521.
- Nakako Y, Fujimoto T, Wada H, et al. Oral metastatic tumor (renal cell carcinoma) in maxillary gingiva: A case report and systematic review. *J Oral Maxillofac Surg Med Pathol.* **2025** Jul;37(4):839-848.
<https://doi.org/10.1016/j.ajoms.2025.01.004>
- National Cancer Institute: Metastasis [internet]. **2025** Jun 07 [cited 2025 Jun 07]. In Ukrainian. Available from:
<https://www.cancer.gov/publications/dictionaries/cancer-terms/def/metastasis#>
- Corominas H, Estrada P, Reina D, Cerdà-Gabaro D. Ultrasonography as a diagnostic tool for skin metastasis of a prostate adenocarcinoma. *Reumatol Clin.* **2016** Jan-Feb;12(1):54-56.
<https://doi.org/10.1016/j.reuma.2015.03.002>
- Martínez-Morán C, Echeverría-García B, Khedaoui R, Borbujo J. Cutaneous metastasis in a patient with lung cancer. *Actas Dermosifiliogr (Engl Ed).* **2018** May;109(4):372-374.
<https://doi.org/10.1016/j.ad.2017.05.019>
- Bai M, Li Z, He P. Ultrasound findings of subcutaneous soft tissue metastasis of renal cell carcinoma: A case report. *Radiol Case Rep.* **2022** Nov 1;18(1):192-195.
<https://doi.org/10.1016/j.radcr.2022.09.055>
- Pickhardt PJ, Pickard RH. Sonography of delayed thyroid metastasis from renal cell carcinoma with jugular vein extension. *AJR Am J Roentgenol.* **2003**;181(1):272-274.
<https://doi.org/10.2214/ajr.181.1.1810272>
- Kanwal R. Metastasis in renal cell carcinoma: Biology and treatment. **2023** Jul;7:100094.
<https://doi.org/10.1016/j.adcanc.2023.100094>

22. Orhan K, Ünsal G. Applications of ultrasonography in maxillofacial/intraoral benign and malignant tumors. In: Ultrasonography in dentomaxillofacial diagnostics. Orhan K, ed. Springer Nature Switzerland AG **2021**; 275-318.
https://doi.org/10.1007/978-3-030-62179-7_18
23. Hirshberg A, Shnaiderman-Shapiro A, Kaplan I, Berger R. Metastatic tumours to the oral cavity - pathogenesis and analysis of 673 cases. *Oral Oncol.* **2008**;44(8):743-752.
<https://doi.org/10.1016/j.oraloncology.2007.09.012>
24. Kelles M, Akarcay M, Kizilay A, Samdanci E. Metastatic renal cell carcinoma to the condyle of the mandible. *J Craniofac Surg.* **2012** Jul;23(4):e302-e303.
<https://doi.org/10.1097/scs.0b013e318252f331>
25. Vasilyeva D, Peters SM, Philipone EM, Yoon AJ. Renal cell carcinoma metastatic to the maxillary gingiva: A case report and review of the literature. *J Oral Maxillofac Pathol.* **2018** Jan;22(Suppl 1):S102-S107.
https://doi.org/10.4103/jomfp.JOMFP_69_17
26. Krushbu, Navik, Sarthak, et al. Unusual metastasis of renal cell carcinoma to upper gingiva: A rare case report. *Int J Life Sci Biotechnol Pharma Res.* **2024** Feb;13(2):42-45.
27. Catalano O, Corvino A. Ultrasound of skin cancer: What we need to know. *Semin Ultrasound CT MR.* **2024**;45(3):216-232.
<https://doi.org/10.1053/j.sult.2023.11.003>
28. Wortsman X. Ultrasound in skin cancer: Why, how, and when to use it? *Cancers.* **2024**; 16(19):3301.
<https://doi.org/10.3390/cancers16193301>
29. Guimarães Ferreira Fonseca L, Bertolizio G, Engelhardt T, Karlsson J. Pediatric point of care airway ultrasound (POCUS): Current evidence and future practice. *Anaesthesiologie.* **2024** Jan 30.
<https://doi.org/10.1007/s00101-024-01377-6>
30. Demidov VH, Cherniak OS, Snisarevskiy PP, et al. Schwannoma of the tongue: ultrasonography. *J Diagn Treat Oral Maxillofac Pathol.* **2022** Nov 30;6(11):138-47.
<https://doi.org/10.23999/j.dcomp.2022.11.2>
31. Fesenko II. Abscess of the left tongue. *J Diagn Treat Oral Maxillofac Pathol.* **2020** Feb 28;4(2):39-40.
<https://doi.org/10.23999/j.dcomp.2020.2.5>
32. Abbott S, Zelesco M, Gibson, D. Tongue sonography: An introduction. *Sonography.* **2020** Aug 03;7(3):118-125.
<https://doi.org/10.1002/sono.12230>
33. Caprioli S, Casaleggio A, Tagliafico AS, et al. High-frequency intraoral ultrasound for preoperative assessment of depth of invasion for early tongue squamous cell carcinoma: Radiological-pathological correlations. *Int J Environ Res Public Health.* **2022** Nov 12;19(22):14900.
<https://doi.org/10.3390/ijerph192214900>
34. Cherniak OS, Fesenko II. A rare odontogenic subperiosteal abscess that involved entire lateral aspect of the mandibular body and medial aspect to the level of mylohyoid ridge: ultrasound examination. *J Diagn Treat Oral Maxillofac Pathol.* **2023** Feb 28;6(2):10-20.
<https://doi.org/10.23999/j.dcomp.2023.2.1>
35. Demidov VH, Ripolovska OV. How multiple the submandibular gland sialoliths can be? *J Diagn Treat Oral Maxillofac Pathol.* **2019** Jul;3(7):174-175.
<https://doi.org/10.23999/j.dcomp.2019.7.2>
36. Dawkins A, Sobieh A, Nair R, Ganesh H. Contemporary sonography of the floor of the mouth: Anatomic highlights, pitfalls, and normal variants. *Contemp Diagn Radiol.* **2022** Mar 15;45(6):1-5.
<https://doi.org/10.1097/01.CDR.0000824016.86045.a5>
37. Rozylo-Kalinowska I, K. Orhan K. General considerations for ultrasound applications in head and neck. In: Ultrasonography in dentomaxillofacial diagnostics. Orhan K, editor. Springer Nature Switzerland AG; **2021**:39-50.
https://doi.org/10.1007/978-3-030-62179-7_3
38. Kim K, Oh D, Shin D, Yoon J. Effectiveness of a hockey-stick probe to localize a catheter fragment in a dog. *J Vet Clin.* **2022** Aug 31;39(4):173-176.
<https://doi.org/10.17555/jvc.2022.39.4.173>
39. Fu L, Chang JJ, Al Hezaimi K, et al. In vivo periodontal ultrasound imaging via a hockey-stick transducer and comparison to periodontal probing: a proof-of-concept study. *Clin Oral Investig.* **2025** Apr 26;29(5):275.
<https://doi.org/10.1007/s00784-025-06346-w>
40. Caglayan F, Bayrakdar IS. The intraoral ultrasonography in dentistry. *Niger J Clin Pract.* **2018** Feb;21(2):125-33.
<http://dx.doi.org/10.4103/1119-3077.197016>
41. Cheng ET, Greene D, Koch RJ. Metastatic renal cell carcinoma to the nose. *Otolaryngol Head Neck Surg.* **2000**;122(3):464.
[https://doi.org/10.1016/s0194-5998\(00\)70069-6](https://doi.org/10.1016/s0194-5998(00)70069-6)
42. Hirshberg A, Berger R, Allon I, Kaplan I. Metastatic tumors to the jaws and mouth. *Head Neck Pathol.* **2014**;8(4):463-474.
<https://doi.org/10.1007/s12105-014-0591-z>
43. Martins FMC, Neto TH, Martins SP, Mendes RB. Metastasis to the oral cavity and jaw bones - A literature review about a case. *Oral Oncol.* **2023**;137:106276.
<https://doi.org/10.1016/j.oraloncology.2022.106276>
44. Küsters B, Westphal JR, Smits D, et al. The pattern of metastasis of human melanoma to the central nervous system is not influenced by integrin alpha(v)beta(3) expression. *Int J Cancer.* **2001** Apr 15;92(2):176-180.
[https://doi.org/10.1002/1097-0215\(200102\)9999:9999%3C::aid-ijc1173%3E3.0.co;2-1](https://doi.org/10.1002/1097-0215(200102)9999:9999%3C::aid-ijc1173%3E3.0.co;2-1)



ЗВІТ ПРО ВИПАДОК/МЕТОДИКА

UKRAINIAN LANGUAGE

Звіт про випадок метастазу світлоклітинної карциноми нирки в ротовій порожнині, що імітував доброякісне новоутворення: можливості та обмеження ультразвукового дослідження

Ольга Сергіївна Черняк^{a,*}, Валентин Григорович Демідов^b, Олександра Андріївна Петриченко^c та В'ячеслав Павлович Близнюк^d

АНОТАЦІЯ

Метастази в ротовій порожнині трапляються рідко, складаючи приблизно 1-1,5% усіх злоякісних новоутворень ротової порожнини. Тому клініцистам важливо напрацьовувати діагностичні дані для ранньої діагностики цього грізного патологічного процесу. Найпоширенішими первинними пухлинами, що метастазують у ротову порожнину, є: легені (21,1%), печінка (12,3%), молочна залоза (10,5%), нирки (10,5%) та колоректальна пухлина (8,8%). Ми сподіваємося, що публікація описаного нижче випадку доповнить міжнародну літературу та введе ультразвукову діагностику на новий рівень. На хірургічне обстеження звернулася 49-річна жінка європеїдної раси зі зростаючим новоутворенням у ротовій порожнині та його грибоподібним прикріпленням до слизової оболонки альвеолярного відростка нижньої щелепи. Новоутворення клінічно імітувало доброякісну патологію. Новоутворення (його передня частина) було досліджено за допомогою ультразвукового дослідження (УЗД) з представленням сірошкальних та кольорових Допплерівських сонограм. Варто відзначити що вестибулярна (передня, менша) частина метастазу в даному випадку не мала характерної ознаки для метастазів в інших ділянках тіла, а саме була відсутня васкуляризація при кольоровому Допплерівському картуванні. У статті також наве-

^a Лікар ультразвукової діагностики; Завідувач відділення ультразвукової діагностики, поліклініка, КНП КОР «Київська обласна клінічна лікарня», Київ, Україна.

^b Лікар -стоматолог-хірург, Центр щелепно-лицевої хірургії КНП КОР «Київська обласна клінічна лікарня» (місце роботи на момент збору матеріалу), Київ, Україна.

^c Лікар-стоматолог-хірург, Центр щелепно-лицевої хірургії КНП КОР «Київська обласна клінічна лікарня» (місце роботи на момент збору матеріалу), Київ, Україна; Власниця, Unique Aesthetic, Батумі, Грузія.

^d Незалежний дослідник: лікар-стоматолог-хірург, Центр щелепно-лицевої хірургії КНП КОР «Київська обласна клінічна лікарня» (місце роботи на момент збору матеріалу), Київ, Україна.

* Кореспонденція: кабінет № 418, поліклініка, КНП КОР «Київська обласна клінічна лікарня», вул. Загорівська, 1, Київ, 04107, Україна. E-mail: cherniak.os@gmail.com (Olha Cherniak). ORCID: <https://orcid.org/0000-0002-6372-0043>.

Тип статті: звіт про випадок/методика.

Для цитування: Cherniak OS, Demidov VH, Petrychenko O, Blyzniuk VP. A case report of oral mucosa metastasis of renal clear cell carcinoma mimicking benign neoplasm: Possibilities and limitations of the ultrasonography. J Diagn Treat Oral Maxillofac Pathol. 2025 Jul;9(7):100307.

На момент публікації даної статті імпакт-фактор (Exaly) журналу становить 0.2. Кожному автору статті нараховується по 20 балів БПР згідно Наказу МОЗ України від 22 лютого 2019 р. № 446. Бібліографічна довідка є підтвердним документом.

Акронім «УЗД» у верхньому правому значку означає, що стаття містить зображення ультразвукової діагностики (УЗД).

Рукопис отриманий 03 червня 2025
Прийнятий до публікації 26 червня 2025
Стаття доступна онлайн 31 липня 2025

<https://doi.org/10.23999/j.dtomp.2025.7.100307>

© 2025 Автор(и). Видано ТОВ «ОМФ ПАБЛІШІНГ». Вебсайт: <https://omfpublishing.com>. Це стаття відкритого доступу під ліцензією CC BY-NC-SA license (<https://creativecommons.org/licenses/by-nc-sa/4.0/>).

дено клінічні дані та дані ортопантомографії. Було проаналізовано обмеження лінійного датчика для повної ультразвукової оцінки новоутворень, розташованих на язичному або, можливо, піднебінному боці альвеолярних відростків. Представлено порівняльний аналіз лінійного ультразвукових датчика та лінійного датчика типу «хокейна ключка». Після проведення аналізу літератури можна стверджувати, що в цій статті вперше представлені дані щодо УЗД метастазів світлоклітинної карциноми нирки на слизовій оболонці рота. Подальше дослідження кожного метастазу в ротовій порожнині за допомогою прямого внутрішньоротового УЗД є надзвичайно важливим для систематизації ультразвукових варіантів цієї патології. У свою чергу, це дозволить клініцистам з більшою ймовірністю встановити правильний попередній діагноз перед біопсією та лікувати цю патологію у спеціалізованих відділеннях.

КЛЮЧОВІ СЛОВА

Метастаз в ротову порожнину, світлоклітинна карцинома нирки, ультразвукове дослідження (УЗД), лінійний датчик, лінійний датчик «хокейна ключка»

U-Impl[®]

SWITZERLAND



№ R3M 804 252 B2

Сертифікат відповідності
технічного регламенту
щодо медичних виробів



Switzerland Aarbergerstrasse 107A, CH-2502
Biel, Phone/Fax +41 323230188
info@u-impl.com
www.u-impl.com